

The right choice, at the right time...for each patient
References

1. Fields, Michael E., et al. How to select patients for lead extraction. Heart Rhythm, Vol 4, Issue 7, July 2007.
2. Voight, Andrew, et al. Rising Rates of Cardiac Rhythm Management Device Infections in the United States: 1996 through 2003. JACC Vol. 48, No. 3, 2006: 590-1.
3. Kleeman Thomas, et al. Annual Rate of Transvenous Defibrillation Lead Defect in Implantable Cardioverter-Defibrillators over a Period of >10 Years. Circulation 2007; 115:2474-2490.
4. Millennium Research Group, Global Markets for Cardiac Rhythm Management Devices 2013
5. Medtronic CRDM Product Performance Report, Mar 2013. Eucomed (2012)
6. Boston Scientific CRM Product Performance Report, Q1 2013. Eucomed (2012)
7. St. Jude Medical CRM Product Performance Report, Apr 2013. Eucomed (2012)
8. Biotronik Product performance Report, July 2013. Kleeman Thomas, et al. Annual Rate of Transvenous Defibrillation Lead Defect in Implantable Cardioverter-Defibrillators over a Period of >10 Years. Circulation 2007; 115:2474-2490
9. Kusumoto et al. 2017 HRS Expert Consensus Statement on Cardiovascular Implantable Electronic Device Lead Management and Extraction. Heart Rhythm, 2017.
10. Pokorney et al. Outcomes Associated with Extraction versus Capping and Abandoning Pacing and Defibrillator Leads Circulation 2017 Oct 10;136(15):1387-1395. doi: 10.1161/ CIRCULATIONAHA.117.027636. Epub 2017 Aug 22.
11. Hussein, Ayman A., et al. "Cardiac Implantable Electronic Device Infections: Added Complexity and Suboptimal Outcomes With Previously Abandoned Leads." JACC: Clinical Electrophysiology (2016).
12. Sohal, M. et al. (2014). Laser lead extraction to facilitate cardiac implantable electronic device upgrade and revision in the presence of central venous obstruction. Europace, 16(1), 81-87.
13. Oginosawa Y, Abe H, Nakashima Y. The incidence and risk factors for venous obstruction after implantation of transvenous pacing leads. Pacing Clin Electrophysiol 2002;25:1605-1611.
14. Kutarski, A., Pietura, R., Młynarczyk, K., Małecka, B., & Główniak, A. (2012). Pacemaker lead extraction and recapture of venous access: technical problems arising from extensive venous obstruction. Cardiology journal, 19(5), 513-517.
15. de Cock CC, et al. Long-term outcome of patients with multiple (> or = 3) noninfected leads: a clinical and echocardiographic study. PACE, Vol 23, No 4, 2000, 423-6
16. Kalin R, Stanton MS. Current clinical issues for MRI scanning of pacemaker and defibrillator patients. PACE. April 2005;28(4):326-328
17. Wilkoff, Bruce, L., et al. Pacemaker Lead Extraction with the Laser Sheath: Results of the Pacing Lead Extraction with Excimer Sheath (PLEXES) Trial. JACC, Vol 33, No. 6, May 1999.
18. Byrd, Charles, et al. Clinical Study of the Laser Sheath for Lead Extraction: The Total Experience in the United States. Journal of Pacing and Electrophysiology. Vol 125, No. 5, May 2002.
19. Wazni, O et. al. Lead Extraction in the Contemporary Setting: The LEXIcon Study: A Multicenter Observational Retrospective Study of Consecutive Laser Lead Extractions, J Am Coll Cardiol, 55:579-586
20. Ryan Azarrafiy, BA; Darren C. Tsang, BS; Bruce L. Wilkoff, MD, FHRS; Roger G. Carrillo, MD, MBA, FHRS. The Endovascular Occlusion Balloon for Treatment of Superior Vena Cava Tears During Transvenous Lead Extraction: A Multi-Year Analysis and An Update to Best Practice Protocol. Circulation: Arrhythmia and Electrophysiology, August 2019
21. Document on file D027562. Bridge can be fully deployed in under one minute (53 seconds) in an animal model when pre-positioned on a guidewire, or in under two minutes (1 minute, 46 seconds) when not pre-positioned.
22. Document on file D027561. When deployed, the Bridge occlusion balloon reduces blood loss by up to 90%, on average, in an animal model of an SVC tear. Testing was conducted in a heparinized porcine model which has shorter SVC length than is typical in humans. A balloon design scaled for use specifically in the porcine model was used in generating this data.
23. Document on file, D026197. In an animal model with SVC tears up to 3.5 cm, with 2 pacing leads and 1ICD lead
24. Elrod, Jodia. Use of Bridge™ Occlusion Balloon in Lead Extraction: Interview with Dr. Roger Carrillo. EP Lab Digest. November, 2016.

25. Wilkoff BL, Love CJ, Byrd CL, Bongiorno MG, Carrillo RG, Crossley GH 3rd, Epstein LM, Friedman RA, Kennergren CE, Mitkowski P, Schaerf RH, Wazni OM; Heart Rhythm Society; American Heart Association. Transvenous lead extraction: Heart Rhythm Society expert consensus on facilities, training, indications, and patient management: this document was endorsed by the American Heart Association (AHA). *Heart Rhythm*. 2009 Jul;6(7):1085-104. doi: 10.1016/j.hrthm.2009.05.020. Epub 2009 May 22. PMID: 19560098.